

Guidelines for psychiatrists for the treatment of alcohol use disorders

Reading and references:

Consensus guidelines:

Agabio R, Sinclair JM, Addolorato G, Aubin H-J, Beraha EM, Caputo F et al (2018) Baclofen for the treatment of alcohol use disorders: the Cagliari statement. *Lancet Psychiatry* 5 (12) 957-960

American Psychiatric Association (2018) Practice guideline for the pharmacological management of patients with alcohol use disorder *American Psychiatric Association*

Haber P, Lintzeris N, Proude E and Lopatko O (2009) Guidelines for the treatment of alcohol problems. *Australian Government Department of Health and Ageing*

Lingford-Hughes AR, Welch S, Peters L and Nutt DJ (2012) Evidence-based guidelines for the pharmacological management of substance abuse, harmful use, addiction and comorbidity: recommendations from the British Association for Psychopharmacology. *Journal of Psychopharmacology* 26 899-952
https://www.bap.org.uk/pdfs/BAP_Guidelines-Addiction.pdf

National Institute for Health and Care Excellence (NICE) UK (2011 – reviewed 2015) Alcohol use disorders: diagnosis, assessment and management of harmful drinking and alcohol dependence.
<https://www.nice.org.uk/guidance/cg115>

National Institute for Health and Care Excellence (NICE) UK (2010 updated 2017) Alcohol use disorders: diagnosis and management of physical complications.
<https://www.nice.org.uk/guidance/cg100>

National Institute for Health and Care Excellence (NICE) UK (2011) Co-existing severe mental illness (psychosis) and substance misuse: assessment and management in healthcare settings.
<https://www.nice.org.uk/guidance/cg120>

National Institute for Health and Care Excellence (NICE) UK (2016) Co-existing severe mental illness: community health and social care services.
<https://www.nice.org.uk/guidance/ng58>

Marel C, Mills KL et al (2016) Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings *Australian Government Department of Health*

Public Health England (2017) Better care for people with co-occurring mental health and alcohol/drug use conditions. *PHE Publications*

Royal College of Psychiatrists (with Royal College of Physicians and Royal College of General Practitioners) (2014) Alcohol and brain damage in adults *Royal College of Psychiatrists, London*

Other references:

Mann K et al (2018) Precision medicine in alcohol dependence: a controlled trial testing pharmacotherapy response among reward and relief drinking phenotypes. *Neuropsychopharmacology* **43** 891-899

Mann K et al (2013) Results of a double-blind placebo-controlled pharmacotherapy trial in alcoholism conducted in Germany and comparison with the US COMBINE study. *Addiction Biology* **18** (6) 937-946

Mark TL, Kassed CA, Vandivort-Warren R, et al: Alcohol and opioid dependence medications: prescription trends, overall and by physician specialty. *Drug Alcohol Depend* **99**(1-3):345-349, 2009 18819759

Pettinati HM, Anton RF, Willenbring ML (2006) The COMBINE Study – an overview of the largest pharmacotherapy study to date for treating alcohol dependence *Psychiatry (Edgmont)* Oct (3) 36-39

Sinclair JM, Chambers SE, Shiles CJ and Baldwin DS (2016) Safety and tolerability of pharmacological treatment of alcohol dependence: comprehensive review of evidence. *Drug Safety* Jul **39**(7) 627-45

Spithoff S (2017) First-line medications for alcohol use disorders among public drug plan beneficiaries in Ontario *Canadian Family Physician* **63**(5) e277-e283

Woolf SH, Grol R, Hutchinson A, Eccles M, Grimshaw J (1999) Potential benefits, limitations and harms of clinical guidelines *British Medical Journal* **318** (7182) 527-530